

GRAPHS OF SCHEMES ASSOCIATED TO GROUP ACTIONS

ALİ ULAŞ ÖZGÜR KİŞİSEL AND ENGİN ÖZKAN

ABSTRACT. Let X be a proper algebraic scheme over an algebraically closed field. We assume that a torus T acts on X such that the action has isolated fixed points. The T -graph of X can be defined using the fixed points and the one dimensional orbits of the T -action. If the upper Borel subgroup of the general linear group with maximal torus T acts on X , then we can define a second graph associated to X , called the A -graph of X . We prove that the A -graph of X is connected if and only if X is connected. We use this result to give a proof of Hartshorne's theorem on the connectedness of Hilbert scheme in the case of d points in $\mathbb{P}^n$.

1. INTRODUCTION

Let X be a proper algebraic scheme over an algebraically closed field k and let $T = (G_m)^n = (k^*)^n$ be an n -dimensional algebraic torus. In the case that X admits a T -action with isolated fixed points one can extract substantial information about X by looking at these fixed points, through equivariant localization or other means (see for instance [5], [7], [11]). In [2] the concept of a T -graph is defined for the Hilbert scheme and its multigraded versions. This definition can be generalized to any setting where the T -action has isolated fixed points. In [9] further properties of T -graphs are obtained and some explicit computations are given.

If X admits an action of a larger group, for instance a Borel subgroup of the general linear group such that the induced maximal torus action has isolated fixed points, then the information that one can extract about X often becomes substantially stronger. As an example, the results of [1] imply that if X admits a compatible (G_m, G_a) -action such that the G_a -fixed locus consists of a single point, then the Poincaré polynomial of X is a product of cyclotomic polynomials.

In this paper, we first generalize the definition of a T -graph to an arbitrary proper algebraic scheme X admitting a T -action with isolated fixed points and show that X is connected if and only if its T -graph is connected. These issues are discussed in section 2. In section 3, assuming that X admits a Borel group action, we define the notion of an A -graph, which is obtained from the fixed locus of X under the action of a principal nilpotent subgroup. The A -graph of X is often substantially smaller compared to the T -graph. For instance, in the setting described in the last sentence of the previous paragraph, the A -graph consists of just a single vertex. We prove that X is connected if and only if its A -graph is connected.

In section 4, we give a proof of Hartshorne's theorem of the connectedness of the Hilbert scheme of $\mathbb{P}^n$ ([10]) in the case where the Hilbert polynomial is a

Key words and phrases. T -graph, A -graph, Hilbert scheme, Borel group action.

constant d , by explicitly computing a spanning subtree of the A -graph of the Hilbert scheme. This proof can be generalized to the case of an arbitrary Hilbert polynomial, but this is not given here. The idea of using Borel fixed monomial ideals is present both in Hartshorne's proof [10] and A. Reeves' proof [13]; the current proof provides a somewhat different organization. Computation of the full A -graphs of Hilbert schemes of points for small values of d and n will be the subject of a future paper.

2. T -GRAPH OF A SCHEME

Let k be an algebraically closed field and X a proper algebraic scheme over k . Let $T = (G_m)^n = (k^*)^n$ be an n -dimensional algebraic torus. Assume that T acts on X such that the fixed point scheme X^T consists of isolated points.

Definition 2.1. *The graph whose vertices correspond to elements of X^T and having an edge between $p, q \in X^T$ if and only if there exists a 1-dimensional T -orbit l whose closure contains both p and q is called the T -graph of X .*

The T -graph of the Hilbert scheme of $\mathbb{P}^n$ was defined in [2]. The T -graphs of multigraded Hilbert schemes of $\mathbb{P}^n$ are also implicitly present in the discussion in [2] although the emphasis there is rather on the induced subgraphs obtained from the ordinary grading. This definition is given in terms of Gröbner degenerations, but it is equivalent to the one above if we assume that the action on this Hilbert scheme is induced from the standard action of the maximal torus of $GL(n+1)$ on $\mathbb{P}^n$. Some details are provided in the two examples below.

Example 2.1. *Let $X = \mathbb{P}^n$ and T be the maximal torus of $GL(n+1)$ containing diagonal matrices $\text{diag}(d_0, \dots, d_n)$ such that $d_i \neq 0$. Consider the standard action of $GL(n+1)$ on X and the corresponding action of T on X given by $\text{diag}(d_0, \dots, d_n) \cdot [x_0 : \dots : x_n] = [d_0 x_0 : \dots : d_n x_n]$. The fixed points of this action are the $n+1$ points of the form $p_i = [0 : \dots : 0 : 1 : 0 : \dots : 0]$ for $i = 0, 1, \dots, n$ where the only nonzero entry of p_i is its i th coordinate. The open subset of the line $l_{ij} = [0 : \dots : \lambda : \dots : \mu : \dots : 0]$, where the only nonzero entries λ, μ are at the i th and j th coordinates respectively, is a 1-dimensional T -orbit. Since p_i and p_j lie in the closure of l_{ij} , any two vertices are connected by an edge. Hence the T -graph of X for this action is the 1-skeleton of the standard n -simplex.*

Example 2.2. *Let $X = \text{Hilb}(\mathbb{P}^n)$, the Hilbert scheme of $\mathbb{P}^n$. Elements of X correspond to homogenous ideals in $k[x_0, x_1, \dots, x_n]$. The T -action on $\mathbb{P}^n$ induces a T -action on X . It is easy to see that the fixed points of this action are precisely the monomial ideals. Indeed, contrary to the claim, assume that the ideal I is a fixed point but it is not a monomial ideal. Then there exists a polynomial $P(x_0, \dots, x_n)$ in I so that none of its monomial summands are in I . Suppose that P is a sum of r monomials. We can assign integer weights $m_0, \dots, m_n$ to the variables $x_0, \dots, x_n$ such that each monomial summand of P has a different weight. Act on P by elements of the form $\text{diag}(t^{m_0}, \dots, t^{m_n})$ for r different values of t . The resulting polynomials $P_1, \dots, P_r$ must be in I because I is a fixed point of the action. Then an elementary computation involving a Van der Monde determinant shows that the set $\{P_1, \dots, P_r\}$ is linearly independent. This implies that each monomial in P can be written as a linear combination of $\{P_1, \dots, P_r\}$,*

contradicting the initial assumption. This contradiction shows that the only fixed points of the T -action are monomial ideals. Thus we determined the vertices of the T -graph. Determining the edges of the T -graph is a more difficult task, see [2] or [9] for a discussion.

Theorem 2.1. *Suppose that T acts on the proper algebraic scheme X with isolated fixed points. Then X is connected if and only if its T -graph is connected.*

Remark 2.1. The analogous statement is false in the category of smooth manifolds. For example, take $k = \mathbb{C}$ and assume that X is S^2 , the 2-sphere. Identify S^2 with the Riemann sphere so that the North pole corresponds to ∞ , the South pole corresponds to 0 and 1 is on the equator. Let $S^1 \times S^1$ act on X in the following way: The first factor acts by rotations parallel to the equator. For the second factor, choose a Möbius transformation M_α sending 0 to 0, ∞ to ∞ and 1 to α where α is not on the equator. Then assume that the action of the second factor is the conjugation of the first action by M_α . Then the fixed points of both actions are the north and south poles. The 1-dimensional orbits of the two actions are two families of circles. A circle in one family intersects circles of the other family transversely on a dense subset. This implies that the $S^1 \times S^1$ -action on X has two fixed points, no 1-dimensional orbits and a single 2-dimensional orbit. We can turn this into a T -action by retracting $T = \mathbb{C}^* \times \mathbb{C}^*$ radially to $S^1 \times S^1$ and composing the retraction with the action described above. The T -graph of X consists in just two disconnected vertices and no edges. Hence the statement is not true in this case.

Lemma 2.1. (i) *There exists a 1-parameter subgroup $G_m \hookrightarrow T$ such that $X^T = X^{G_m}$.*

(ii) *There exists a 2-parameter subgroup $G_m \times G_m \hookrightarrow T$ such that the T -graph of X agrees with the $G_m \times G_m$ -graph of X .*

Proof It is enough to prove both statements in the case where X is irreducible.

(i) By the Bialynicki-Birula decomposition [3] there exists a big cell $U \subset X$ that contains a unique fixed point a and the T -action restricted to U is conjugate to a linear action of T on a vector space V . Since the action has a unique fixed point, all weights of this action must be nontrivial. Suppose that we choose a basis that diagonalizes the action, namely $t = (t_1, \dots, t_n) \in T$ acts on V by $\text{diag}(\lambda_1(t), \dots, \lambda_N(t))$. Assume that $\lambda_i(t) = t_1^{e_{i1}} t_2^{e_{i2}} \dots t_n^{e_{in}}$. Since there are no trivial weights, for each i there exists j such that $e_{ij} \neq 0$. Now consider an arbitrary one-parameter subgroup $\varphi : G_m \hookrightarrow T$ so that $\varphi(s) = (s^{a_1}, s^{a_2}, \dots, s^{a_n})$. The weights restricted to this subgroup are nontrivial if and only if for each i , the sum $\sum_{j=1}^n e_{ij} a_j$ is nonzero. Since this is a set of finitely many nontrivial inequalities in integers, there exists $(a_1, \dots, a_n)$ such that $U^T = U^{G_m}$. Now replace X by $X - U$ and proceed inductively. Since at every step finitely many equalities are to be avoided and there are finitely many steps, there exist one-parameter subgroups $\varphi : G_m \hookrightarrow T$ such that $X^{G_m} = X^T$.

(ii) Let U and V be as in part (i) of the proof. The T -graph of X has a unique vertex v on U corresponding to the unique fixed point. The edges of the graph emanating from v correspond to lines spanned by eigenvectors of the T -action on V . Choose a basis as in part (i). A two parameter subgroup

$\varphi : G_m \times G_m \hookrightarrow T$ induces the same edges as T if and only if no new vertices are produced and the eigenspaces of the $G_m \times G_m$ -action are the same as the eigenspaces of the T -action. Say $\varphi(s, u) = (s^{a_1}u^{b_1}, s^{a_2}u^{b_2}, \dots, s^{a_n}u^{b_n})$. Then $\lambda_i(s, u) = s^{\sum_{j=1}^n a_j e_{ij}} u^{\sum_{j=1}^n b_j e_{ij}}$. Therefore the T -graph restricted to U agrees with the $G_m \times G_m$ -graph restricted to U if and only if the following inequalities are satisfied for all pairs i, k with $(e_{i1}, \dots, e_{in}) \neq (e_{k1}, \dots, e_{kn})$:

$$\left(\sum_{j=1}^n a_j e_{ij}, \sum_{j=1}^n b_j e_{ij} \right) \neq \left(\sum_{j=1}^n a_j e_{kj}, \sum_{j=1}^n b_j e_{kj} \right)$$

and $(\sum_{j=1}^n a_j e_{ij}, \sum_{j=1}^n b_j e_{ij}) \neq (0, 0)$ for all i . Since a finite set of nontrivial equalities need to be avoided, there exist solutions for a_i, b_i . The rest of the argument proceeds as in (i), by considering $X - U$ and doing induction. $\square$

We will adapt the strategy in [4] to our case in order to prove theorem 2.1. We now recall a definition from [4] :

Definition 2.2. *Let A, X_1, X_2 be schemes. We say that X_1 is simply A -equivalent to X_2 if there exists an isomorphism $X_1 - X_2 \rightarrow A \times Y$ for some scheme Y . The equivalence closure of simple A -equivalence is called A -equivalence.*

Let K be a finite field so that $\text{char}(K) \neq \text{char}(k)$. Let $H^i(X)$ denote the i -th étale cohomology group of X with proper supports, with coefficients in the constant sheaf K .

Lemma 2.2. *Say X_1 and X_2 are schemes such that X_1 is $G_m \times G_m$ -equivalent to X_2 . Then $H^0(X_1) \cong H^0(X_2)$.*

Proof Since $H^0(G_m) = 0$ and $H^1(G_m) = K$, by the Künneth formula we obtain $H^0(G_m \times G_m) = H^1(G_m \times G_m) = 0$.

It is enough to prove the statement when X_1 is simply $G_m \times G_m$ -equivalent to X_2 . Hence we may assume that $X_1 - X_2 \cong U = G_m \times G_m \times U_1$. If U_1 is singular then we can stratify its singular locus and break this equivalence into finitely many steps in each of which U_1 is nonsingular. Hence we may assume that U_1 is nonsingular. By the Künneth formula, there exists a spectral sequence such that $E_2^{p,q} = H^p(U_1) \otimes H^q(G_m \times G_m)$ converging to $H^{p+q}(U)$.

First, let us consider the case of $\dim(U) = 2$. Then U is isomorphic to a disjoint union of copies of $G_m \times G_m$. It suffices to consider one copy at a time, so assume that $U \cong G_m \times G_m$. If $\bar{U} = U$ then $H^0(\bar{U}) = H^0(U) = 0$. The excision sequence in cohomology says that $0 \rightarrow H^0(X_1 - \bar{U}) \rightarrow H^0(X_1) \rightarrow H^0(\bar{U})$ is exact. But $X_1 - \bar{U} = X_2$ and $H^0(\bar{U}) = 0$. Therefore $H^0(X_1) \cong H^0(X_2)$. If $\bar{U} \neq U$ then first consider the excision sequence with respect to the inclusion $U \hookrightarrow \bar{U}$. Then

$$H^0(U) \rightarrow H^0(\bar{U}) \rightarrow H^0(\bar{U} - U) \rightarrow H^1(U)$$

is exact. But $H^0(U) = H^1(U) = 0$, therefore $H^0(\bar{U}) \cong H^0(\bar{U} - U)$. Now, consider the excision sequences with respect to the inclusions $\bar{U} \hookrightarrow X_1$ and $\bar{U} - U \hookrightarrow X_1 - U$.

$$\begin{array}{ccccccc}
 0 & \longrightarrow & H^0(X_1 - \bar{U}) & \longrightarrow & H^0(X_1) & \longrightarrow & H^0(\bar{U}) \longrightarrow H^1(X_1 - \bar{U}) \\
 & & \downarrow \cong & & \downarrow \alpha & & \downarrow \cong \\
 0 & \longrightarrow & H^0(X_1 - \bar{U}) & \longrightarrow & H^0(X_1 - U) & \longrightarrow & H^0(\bar{U} - U) \longrightarrow H^1(X_1 - \bar{U})
 \end{array}$$

By the 5-lemma, the map $\alpha : H^0(X_1) \rightarrow H^0(X_1 - U) = H^0(X_2)$ is an isomorphism, which finishes the proof in this case.

Next, consider the case of $\dim(U) > 2$. We may assume that none of the connected components of U_1 is complete. Else, stratify U_1 and subdivide the equivalence into more steps, where the stratum with $\dim(U_1) = 0$ is taken care of as above. Hence $H^0(U_1) = 0$. Since $E_2^{p,q}(U) = H^p(U_1) \otimes H^q(G_m \times G_m)$ we get $E_2^{0,0}(U) = E_2^{0,1}(U) = E_2^{1,0}(U) = 0$. This implies $H^0(U) = H^1(U) = 0$. The rest of the argument is identical to the one in the first case above. $\square$

Proof of Theorem 2.1 Let Y denote the union of 0 and 1-dimensional orbits of the T -action on X . By lemma 2.1 part (ii), there exists a subgroup of T isomorphic to $G_m \times G_m$ whose 0 and 1-dimensional orbits agree with those of T . It is clear that X is $G_m \times G_m$ -equivalent to Y . Hence, by lemma 2.2, $H^0(X) \cong H^0(Y)$. Since X is proper, X is connected if and only if $H^0(X) \cong K$ and the same holds for Y . On the other hand, by definition, Y is connected if and only if the T -graph of X is connected. This establishes the claim. $\square$

3. ADDITIVE GROUP ACTIONS AND A -GRAPHS

Let G_a denote the 1-dimensional connected additive algebraic group.

Definition 3.1. Let X be an algebraic scheme over k having G_m and G_a -actions such that

$$\begin{aligned}
 \lambda : G_m \times X &\rightarrow X, & ((t, x) &\mapsto \lambda(t) \cdot x) \\
 \theta : G_a \times X &\rightarrow X, & ((u, x) &\mapsto \theta(u) \cdot x)
 \end{aligned}$$

The actions λ and θ are said to be compatible if there exists an integer $p \geq 1$ such that

$$\lambda(t) \cdot \theta(u) \cdot \lambda(t^{-1}) = \theta(t^p \cdot u)$$

for all $t \in G_m$ and $u \in G_a$.

Suppose that X admits compatible G_m and G_a -actions. Then the G_a -fixed locus X^{G_a} of X is G_m -invariant. Indeed, suppose that $x \in X^{G_a}$. Then $\theta(u)\lambda(t)x = \lambda(t)\theta(t^{-p}u)x = \lambda(t)x$. Thus $\lambda(t)x \in X^{G_a}$.

Let $E_{i,j}$ denote the elementary matrix whose ij -entry is 1 and all others 0. Say $E = \sum_{i=1}^{n-1} E_{i,i+1}$. Then a copy of G_a consisting in matrices of the form $\exp(aE)$ for $a \in k$ sits inside $GL(n)$. Let T be the maximal torus of $GL(n)$ consisting of diagonal matrices. Then it is easy to check that 1-parameter subgroups G_m of T compatible with this G_a are subgroups containing matrices of the form:

$$\text{diag}(t^c, t^{c-p}, t^{c-2p}, \dots, t^{c-(n-1)p}).$$

Say X is a proper algebraic scheme over k . Let B be the upper Borel subgroup of $GL(n)$ and T the maximal torus in B consisting of diagonal matrices. Suppose

that B acts on X such that the induced action of T on X has isolated fixed points. Let G_a be the subgroup of B consisting in the matrices $\exp(aE)$ as above. The following lemma is a refinement of lemma 2.1:

Lemma 3.1. *There exists a 2-parameter subgroup $G_m \times G_m \hookrightarrow T$ such that*

- (i) *the T -graph of X agrees with the $G_m \times G_m$ -graph of X ,*
- (ii) *actions of both G_m factors are compatible with the G_a -action.*

Proof In order for condition (ii) to be satisfied, the two G_m factors should contain matrices of the form $\text{diag}(t^{a_1}, \dots, t^{a_n})$ and $\text{diag}(t^{b_1}, \dots, t^{b_n})$ respectively where $a_i = c_1 - (i-1)p_1$ and $b_i = c_2 - (i-1)p_2$ for some c_1, c_2, p_1, p_2 . One can check that the inequalities $(\sum_{j=1}^n a_j e_{ij}, \sum_{j=1}^n b_j e_{ij}) \neq (\sum_{j=1}^n a_j e_{kj}, \sum_{j=1}^n b_j e_{kj})$ and $(\sum_{j=1}^n a_j e_{ij}, \sum_{j=1}^n b_j e_{ij}) \neq (0, 0)$ in the proof of lemma 2.1 are still nontrivial. Therefore there exist choices of c_1, c_2, p_1, p_2 such that all inequalities are satisfied. $\square$

Definition 3.2. *Suppose that X is a proper scheme admitting a Borel group action and let $G_m \times G_m$ and G_a be as in 3.1. Then the $G_m \times G_m$ graph of X^{G_a} is called the A -graph of X associated to this data.*

Recall Horrocks' theorem [10] saying that the fixed locus X^S of the action of a solvable group S on a proper scheme X is connected if and only if X itself is connected. This holds in particular for G_a -actions.

Lemma 3.2. *The A -graph of X is independent of the choice of the $G_m \times G_m$ in its definition. Its vertices are the vertices of the T -graph which are G_a -fixed and its edges are the edges of the T -graph which are G_a -invariant.*

Proof The vertices of the A -graph are $G_m \times G_m$ -fixed points in X^{G_a} . By the choice of $G_m \times G_m$, these are precisely the T -fixed points in X which are also G_a -fixed. Similarly, 1-dimensional T -orbits agree with 1-dimensional $G_m \times G_m$ -orbits. Say l is such an orbit whose closure contains the fixed points P and Q . Suppose that l is G_a -invariant. But in this case the closure of l , which is isomorphic to $\mathbb{P}^1$ admits a G_a -action with at least two fixed points P and Q . By Horrocks' theorem, the G_a -fixed locus of this action on $\mathbb{P}^1$ must be connected, therefore all of l must lie in the G_a -fixed locus X^{G_a} . Therefore the edges of the A -graph are precisely the 1-dimensional T -orbits which are G_a -invariant or equivalently G_a -fixed. $\square$

Theorem 3.1. *Suppose that the Borel subgroup B of $GL(n)$ acts on the proper algebraic scheme X such that the induced T -action has isolated fixed points. Then the A -graph of X is connected if and only if X is connected.*

Proof By Horrocks' theorem, X is connected if and only if X^{G_a} is connected. Applying theorem 2.1 to X^{G_a} we see that X^{G_a} is connected if and only if the A -graph of X is connected. The result follows. $\square$

4. A-GRAPH OF $\text{Hilb}^d(\mathbb{P}^n)$

In this section we will show that the A -graph of $\text{Hilb}^d(\mathbb{P}^n)$ is connected for any $d > 0$ and $n > 0$. Given theorem 3.1, this will provide us with a proof of Hartshorne's theorem on the connectedness of the Hilbert scheme [8] in the

restricted case of a constant Hilbert polynomial. Instead of determining the A -graph completely, we will explicitly describe the vertices and a spanning subtree of this graph. The case of $n = 2$ was treated in [12]. The arguments here can be generalized to any Hilbert polynomial $p(t)$, but we do not carry this out here.

Consider the standard representation of $GL(n+1)$ on K^{n+1} by matrix multiplication. This induces a $GL(n+1)$ -action on $\mathbb{P}^n(K)$ and consequently the upper Borel subgroup B of $GL(n+1)$ acts on $\mathbb{P}^n$. Selecting the subgroup G_a of B as matrices of the form $\exp(aE)$ as in the previous section, the induced G_a -action θ on $\mathbb{P}^n$ is given by

$$\theta(a)([X_0 : \dots : X_n]) = [X_0 + aX_1 + \dots + \frac{a^n}{n!}X_n : X_1 + aX_2 + \dots + \frac{a^{n-1}}{(n-1)!}X_n : \dots : X_n].$$

In particular, the G_a -action on $\mathbb{P}^n$ has a unique fixed point $[1 : 0 : \dots : 0]$.

Consider the induced B -action on $K[X_0, \dots, X_n]$. It takes homogenous ideals to homogenous ideals and the action is flat. Therefore, there is an induced B -action on the Hilbert scheme $\text{Hilb}^d(\mathbb{P}^n)$.

Let us first determine the vertices of the A -graph of $\text{Hilb}^d(\mathbb{P}^n)$. Any vertex must be T -fixed, therefore it must be a monomial ideal. The G_a -fixed monomial ideals agree with the Borel fixed ideals ([6], thm. 15.23):

Lemma 4.1. *A homogenous monomial ideal I is G_a -fixed if and only if it has the following property: For every $f \in I$ divisible by X_i one has $\frac{X_{i+1}}{X_i}f \in I$.*

Proof Suppose that I satisfies the given condition. Since $\theta(a)(X_i) = X_i + aX_{i+1} + \dots + \frac{a^{n-i}}{(n-i)!}X_n$ it is clear that if m is a monomial in I then every monomial summand of $\theta(a)m$ remains in I . Therefore I is G_a -fixed. Conversely, suppose that I is G_a -fixed. Say $f \in I$ and $X_i|f$. Without loss of generality, we may assume that $f = \prod X_j^{e_j}$ is a monomial. Then $\theta(a)(f) = \prod (X_j + aX_{j+1} + \dots)^{e_j} \in I$. Since $e_i \geq 1$, by binomial expansion the monomial $\frac{X_{i+1}}{X_i}f$ appears among the monomial summands of $\theta(a)(f)$. Since I is a monomial ideal, $\frac{X_{i+1}}{X_i}f \in I$. $\square$

Let us now pass to the affine chart $X_0 \neq 0$, which contains the fixed point $[1 : 0 : \dots : 0]$. Say $x_i = \frac{X_i}{X_0}$ be affine coordinates on this chart. Since the support of a G_a -fixed monomial ideal in the Hilbert scheme must be the point $[1 : 0 : \dots : 0]$, without loss of generality we may consider monomial ideals in the variables $x_1, \dots, x_n$ from now on.

Definition 4.1. *Suppose that I and J are two G_a -fixed monomial ideals. Say $m_1, m_2, \dots, m_k$ and $m'_1, \dots, m'_k$ are monomials such that $m_i < m'_i$ in lexicographic order for each i . Assume that each m_i and m'_i/x_1 belongs to the minimal set of generators of I . We say that $(m_1, \dots, m_k) \rightarrow (m'_1, \dots, m'_k)$ is a move from I to J if J can be obtained from I by replacing m_i by m_i/x_1 and m'_i/x_1 by $m'_i x_j/x_1$ for $j = 1, \dots, n$ in the minimal set of generators of I .*

Remark 4.1. After a move, the set of generators obtained for J need not be minimal anymore.

Remark 4.2. A move from I to J can easily be expressed in terms of socles of these ideals: The move $(m_1, \dots, m_k) \rightarrow (m'_1, \dots, m'_k)$ removes m_i/x_1 from the

socle of I and adds m'_i/x_1 to it for each $i = 1, \dots, k$. Such an operation does not change the number of elements in the socle. Therefore, if there exists a move from I to J then I and J must have the same Hilbert polynomial.

Example 4.1. Say $n = 3$. Suppose that

$$I = \langle x_1^3, x_1^2x_2, x_1^2x_3, x_1x_2^2, x_1x_2x_3, x_1x_3^2, x_2^3, x_2^2x_3, x_2x_3^2, x_3^3 \rangle$$

$$J = \langle x_1^4, x_1^2x_2, x_1x_2^2, x_1^2x_3, x_1x_2x_3, x_2^3, x_2^2x_3, x_3^3 \rangle.$$

Take $m_1 = x_1x_3^2$ and $m'_1 = x_1^4$. Then J can be obtained from I by the move $m_1 \rightarrow m'_1$.

Definition 4.2. Say $m = x_1^{a_1}x_2^{a_2} \dots x_n^{a_n}$ is a monomial. Define the weight of m to be

$$w(m) = \sum_{j=1}^n a_j(n-j)$$

The weight of a monomial ideal I is defined to be the sum of the weights of all monomials in the socle of I .

Lemma 4.2. Suppose that I and J are two G_a -fixed monomial ideals. If there exists a move from I to J then $w(J) > w(I)$.

Proof By remark 4.2, it suffices to show that $w(m_i/x_1) < w(m'_i/x_1)$, namely $w(m_i) < w(m'_i)$. Since $m_i < m'_i$ in lexicographic ordering and I is G_a -fixed, $\deg(m_i) \leq \deg(m'_i/x_n) < \deg(m'_i)$. The result follows. $\square$

Let I be a G_a -fixed monomial ideal minimally generated by $S = \{m_1, m_2, \dots, m_k\}$. Let $S^{(d)}$ denote the subset of degree d monomials in S .

Definition 4.3. Let d be the largest integer such that $S^{(d)} \neq \emptyset$ and $S^{(d)}$ contains a monomial other than x_1^d . Let j be the maximal positive integer such that x_j divides some element of $S^{(d)}$. Let l be the maximal integer such that x_j^l divides at least two elements of $S^{(d)}$. Let

$$\hat{S} = \{m_i \in S^{(d)} : x_j^l | m_i, x_1 | m_i\}$$

Let us write elements of $\hat{S}$ in increasing order with respect to lex. Namely, let $\hat{S} = \{m_1, m_2, \dots, m_r\}$ such that $m_1 < m_2 < \dots < m_r$.

Lemma 4.3. Suppose that I , j , l and $\hat{S} = \{m_1, \dots, m_r\}$ are as above and that $r \geq 2$. Then there exists a positive integer s such that the ideal J obtained from I by the move

$$(m_1, m_2, \dots, m_s) \mapsto \left(\left(\frac{x_1}{x_j}\right)^l x_1 m_1, \left(\frac{x_1}{x_j}\right)^l x_1 m_2, \dots, \left(\frac{x_1}{x_j}\right)^l x_1 m_s\right)$$

is also G_a -fixed.

Proof Since the largest element in $\hat{S}$ is $x_1^{d-l}x_j^l$ and $r \geq 2$, we have $m_1 \neq x_1^{d-l}x_j^l$. Then m_1 must be divisible by x_i for some $1 < i < j$. By definition of $\hat{S}$, the monomial $\left(\frac{x_1}{x_j}\right)^l m_1$ must be in I , hence the move $m_1 \mapsto x_1 \left(\frac{x_1}{x_j}\right)^l m_1$ is valid. If the resulting ideal is not G_a -fixed then $\frac{x_1}{x_i} \left(\frac{x_1}{x_j}\right)^l m_1$ must be in I for some $1 < i < j$.

Select the maximal value of i for which this holds. Then by G_a invariance of I , the monomial $\frac{x_1}{x_i}m_1 = m_k$ must be in $\hat{S}$. Apply the move

$$(m_1, m_2, \dots, m_k) \mapsto \left(\left(\frac{x_1}{x_j} \right)^l x_1 m_1, \left(\frac{x_1}{x_j} \right)^l x_1 m_2, \dots, \left(\frac{x_1}{x_j} \right)^l x_1 m_k \right).$$

Repeat the same argument with m_1 replaced by m_k . After finitely many steps, the process terminates. $\square$

Lemma 4.4. *Suppose that I, j, l and $\hat{S}$ are as above but $r = 1$, so that $\hat{S} = \{m_1\}$. Then there exists $h \leq l$ and $k \geq h$ such that the ideal J obtained from I by the move*

$$m_1 \mapsto \left(\frac{x_1^k}{x_j^h} \right) x_1 m_1$$

is also G_a -fixed. If $h < l$ then $k = h$.

Proof First suppose that $l \geq 2$. If the monomial $\frac{x_1^l}{x_j^l}m_1$ is in I then the move

$$m_1 \mapsto \left(\frac{x_1^l}{x_j^l} \right) x_1 m_1$$

is valid and the resulting ideal is G_a -fixed. If not, then by definition of $\hat{S}$, the monomial $\frac{x_1^{l-1}}{x_j^{l-1}}m_1$ is in I so the move

$$m_1 \mapsto \left(\frac{x_1^{l-1}}{x_j^{l-1}} \right) x_1 m_1$$

is valid and again the resulting ideal is G_a -fixed.

It remains to consider the case $l = 1$. Then $m_1 = x_1^{d-l}x_j^l$. Suppose that d' is minimal such that $x_1^{d'} \in I$. Then the move

$$m_1 \mapsto \left(\frac{x_1^{d'-d+l}}{x_j^l} \right) x_1 m_1$$

is valid and again, the ideal obtained is G_a -fixed. $\square$

Lemmas 4.2, 4.3 and 4.4 imply that starting from any G_a -fixed monomial ideal I there exists a finite sequence of moves through the vertices of the A -graph of the Hilbert scheme ending at the ideal $\langle x_1^d, x_2, \dots, x_{n-1}, x_n \rangle$. Next we want to show that if we can pass from vertex v to vertex w by a move as described above, then these two vertices are connected with an edge of the A -graph.

Lemma 4.5. *Suppose that I is minimally generated by $m_1, m_2, \dots, m_k$ and $\hat{S} = \{m_1, \dots, m_r\}$, s, J, j and l are as in lemma 4.3. Assume that J is obtained from I by the move*

$$(m_1, m_2, \dots, m_s) \mapsto (m'_1, m'_2, \dots, m'_s).$$

Reorder the minimal generating set for I such that the first s elements are $m'_1/x_1, \dots, m'_s/x_1$ and the remaining ones are $\hat{m}_{s+1}, \dots, \hat{m}_k$. Suppose that $x_1^{a_i} | m_i$ but $x_1^{a_i+1}$ does not divide m_i .

Then,

(i) *The family of ideals*

$$I_t = \left\langle \frac{m'_1}{x_1} + c_1 t \frac{m_1}{x_1}, \frac{m'_2}{x_1} + c_2 t \frac{m_2}{x_1}, \dots, \frac{m'_s}{x_1} + c_s t \frac{m_s}{x_1}, \hat{m}_{s+1}, \dots, \hat{m}_k \right\rangle$$

where $c_i = \frac{(a_i + l)!}{(a_i - 1)!}$ forms an edge of the A -graph.

(ii) $I_0 = I$.

(iii) $\lim_{t \rightarrow \infty} I_t = J$.

Proof Assuming that I_t is defined as in (i), the proof of (ii) immediately follows by putting $t = 0$.

In order to prove (iii) it suffices to show that $m'_i \in J$ and $\frac{m_i}{x_1} \in J$ for each $i \in \{1, \dots, s\}$. Since $m_i \in I_t$ for each t , the first of these follows from the equality

$$m'_i = x_1 \left(\frac{m'_i}{x_1} + c_i t \frac{m_i}{x_1} \right) - c_i t m_i.$$

The second can be shown by dividing $\frac{m'_i}{x_1} + c_i t \frac{m_i}{x_1}$ by $c_i t$ and taking t to ∞ .

Now let us prove (i). We need to show that I_t is G_a -fixed for each t . Let us write the generators $\{\hat{m}_{s+1}, \dots, \hat{m}_k\}$ in decreasing order. The last monomial, which is of the form x_n^p , is G_a -fixed. The G_a -action on a monomial m produces monomials less than or equal to m in the lexicographic order. Using this fact and descending induction on the index, we see that $\langle \hat{m}_{s+1}, \dots, \hat{m}_k \rangle$ is G_a -fixed. Similarly we see that $\langle \frac{m'_1}{x_1} + c_1 t \frac{m_1}{x_1}, \hat{m}_{s+1}, \dots, \hat{m}_k \rangle$ is G_a -fixed. We want to show by induction that each segment $\langle \frac{m'_1}{x_1} + c_1 t \frac{m_1}{x_1}, \frac{m'_2}{x_1} + c_2 t \frac{m_2}{x_1}, \dots, \frac{m'_i}{x_1} + c_i t \frac{m_i}{x_1}, \hat{m}_{s+1}, \dots, \hat{m}_k \rangle$ is G_a -fixed. In order to complete the induction step, we need to see that the G_a -action on $\frac{m'_i}{x_1} + c_i t \frac{m_i}{x_1}$ yields the monomials $\frac{m'_r}{x_r}$ and $\frac{m_r}{x_r}$ in a term proportional to $\frac{m'_r}{x_1} + c_r t \frac{m_r}{x_1}$. First assume $a_i = a_r$, so that $c_i = c_r$. Then the coefficient of $\frac{m_r}{x_1}$ resulting from the G_a -action on $\frac{m_i}{x_1}$ is a weighted sum over all decreasing sequences (in lexicographic order) of monomials starting from $\frac{m_i}{x_1}$ and ending at $\frac{m_r}{x_1}$. Since the move $m_i \rightarrow m'_i$ does not change any exponents other than x_1 and x_j , the same set of decreasing sequences occurs from $\frac{m'_i}{x_1}$ to $\frac{m'_r}{x_r}$. Therefore the aforementioned claim holds. On the other hand if $a_i \neq a_r$, the only difference in the sequences is in the steps transferring powers of x_1 to the other variables, and the choice of c_i balances the count. $\square$

Corollary 4.1. *The ideals in lemma 4.1 correspond to the vertices and the families of ideals in lemma 4.5 correspond to edges of a spanning subtree of the A -graph of $\text{Hilb}^d(\mathbb{P}^n)$. In particular, this graph is connected, hence by theorem 3.1, $\text{Hilb}^d(\mathbb{P}^n)$ is connected.*

REFERENCES

- [1] Akyıldız, E. , Carrell, J. B. *Cohomology of projective varieties with regular SL_2 -actions*, Manuscripta Math. , 58 (1987), 473–486 [1](#)
- [2] Altmann, K. , Sturmfels, B. *The graph of monomial ideals*, J. Pure Appl. Algebra, 201(1–3) (2005), 250–263 [1](#), [2](#), [2.2](#)
- [3] Białyński-Birula, A. *Some theorems on actions of algebraic groups*, Ann. of Math. (2) 98 (1973), 480–497 [2](#)

- [4] Biaynicki-Birula, A. *On fixed point schemes of actions of multiplicative and additive groups*, Topology 12 (1973), 99-103 [2](#)
- [5] Bialynicki-Birula, A. , Carrell, J. B. , McGovern, W. M. *Algebraic quotients. Torus actions and cohomology. The adjoint representation and the adjoint action*, volume 131 of Encyclopaedia of Mathematical Sciences. Springer-Verlag, Berlin, 2002. Invariant Theory and Algebraic Transformation Groups, II. [1](#)
- [6] Eisenbud, D. *Commutative Algebra with a View Towards Algebraic Geometry*, (1995), Springer-Verlag, New York [4](#)
- [7] Goresky, M. , Kottwitz, R. , MacPherson, R. *Equivariant cohomology, Koszul duality, and the localization theorem* Invent. Math., 131(1) (1998), 25-83 [1](#)
- [8] Hartshorne, R. *Connectedness of the Hilbert Scheme*, Inst.Haute Etude Sci. Pub.Math. 29 (1966), 261-309 [4](#)
- [9] Hering, M. , MacLagan, D. *The T-graph of a multigraded Hilbert scheme*, Experiment. Math. Volume 21, Issue 3 (2012), 280–297 [1](#), [2.2](#)
- [10] Horrocks, G. *Fixed point schemes of additive group schemes*, Topology 8 (1969), 233–242 [1](#), [3](#)
- [11] Iversen, B. *A fixed point formula for action of tori on algebraic varieties* Invent. Math., 16 (1972), 229-236 [1](#)
- [12] Özkan, E. *Fixed point scheme of the Hilbert scheme under a 1-dimensional additive algebraic group action* Ph. D. Thesis, Middle East Technical University, 2011 [4](#)
- [13] Reeves, A. A. *The radius of the Hilbert scheme* J. of Algebraic Geometry, 4 (1995), 639-657 [1](#)

MIDDLE EAST TECHNICAL UNIVERSITY, 06531 ANKARA, TURKEY

E-mail address: `akisisel@metu.edu.tr`

SÜLEYMAN DEMIREL UNIVERSITY, ISPARTA, TURKEY

E-mail address: `enginozkan@sdu.edu.tr`